

Composition-Diamond lemma for λ -differential associative algebras with multiple operators*

Jianjun Qiu

Mathematics and Computational Science School

Zhanjiang Normal University

Zhanjiang 524048, China

jianjunqiu@126.com

Yuqun Chen

School of Mathematical Sciences, South China Normal University

Guangzhou 510631, China

yqchen@scnu.edu.cn

Abstract: In this paper, we establish the Composition-Diamond lemma for λ -differential associative algebras over a field K with multiple operators. As applications, we obtain Gröbner-Shirshov bases of free λ -differential Rota-Baxter algebras. In particular, linear bases of free λ -differential Rota-Baxter algebras are obtained and consequently, the free λ -differential Rota-Baxter algebras are constructed by words.

Key words: λ -differential Rota-Baxter algebra, λ -differential associative algebra with multiple operators, Gröbner-Shirshov basis.

AMS 2000 Subject Classification: 16S15, 13P10, 16W99, 17A50

1 Introduction

Let K be a field and $\lambda \in K$. A differential algebra of weight λ , namely, a λ -differential algebra (see [15, 16]), is an associative K -algebra R with a λ -differential operator $D : R \rightarrow R$ such that

$$D(xy) = D(x)y + xD(y) + \lambda D(x)D(y), \forall x, y \in R.$$

A Rota-Baxter algebra of weight λ (see [2, 19]) is an associative K -algebra R with a Rota-Baxter operator $P : R \rightarrow R$ such that

$$P(x)P(y) = P(xP(y)) + P(P(x)y) + \lambda P(xy), \forall x, y \in R.$$

*Supported by the NNSF of China (No.10771077) and the NSF of Guangdong Province (No.06025062).

Similar to the relation of integral and differential operators, L. Guo and W. Keigher [15] introduced the notion of λ -differential Rota-Baxter algebra which is a K -algebra R with a λ -differential operator D and a Rota-Baxter operator P such that $DP = Id_R$.

There have been some constructions of free Rota-Baxter algebras (commutative and associative). We note that G.-C. Rota [19] and P. Cartier [8] gave the explicit constructions of the free commutative Rota-Baxter algebras on a set when $\lambda = 1$, namely, the shuffle Baxter and standard Baxter algebras, respectively. Recently, L. Guo and W. Keigher [13, 14] constructed the free commutative Rota-Baxter algebras (with identity or without identity) for any $\lambda \in K$ by the mixable shuffle product. These algebras are now called the mixable shuffle product algebras. In fact, these algebras generalize the classical construction of shuffle product algebras. K. Ebrahimi-Fard and L. Guo [11] constructed recently the free associative Rota-Baxter algebras by Rota-Baxter words.

E. Kolchin [16] considered the differential algebra and constructed a free differential algebra. L. Guo and W. Keigher [15] dealt with a generalization of this algebra. Also in [15], the free λ -differential Rota-Baxter algebra was obtained by using the free Rota-Baxter algebra on planar decorated rooted trees.

K. Ebrahimi-Fard and L. Guo [12] used rooted trees and forests to give an explicit construction of free noncommutative Rota-Baxter algebras on modules and sets. K. Ebrahimi-Fard, J. M. Gracia-Bondia and F. Patras [10] gave the solution of the word problem for free non-commutative Rota-Baxter algebra. A free Rota-Baxter algebra was constructed on decorated trees by M. Aguiar and M. Moreira [1].

The concept of multi-operators algebras (Ω -algebras) was first introduced by A. G. Kurosh in [17, 18]. Also, Kurosh noticed that free Ω -algebras share many of the combinatorial properties of free non-associative algebras. On the other hand, the Gröbner-Shirshov bases theory for Lie algebras was first considered by A. I. Shirshov [20]. In fact, Shirshov [20] defined the composition of two Lie polynomials and established the Composition lemma for the Lie algebras. Later on, L. A. Bokut [4] specialized the approach of Shirshov to associative algebras, see also G. M. Bergman [3]. For commutative polynomials, this lemma is known as the Buchberger's Theorem in [6, 7].

Gröbner-Shirshov bases for Ω -algebras were given in the paper of V. Drensky and R. Holtkamp [9]. In a recent paper of L. B. Bokut, Y. Chen and J. Qiu [5], the Composition-Diamond lemma is established for associative Ω -algebras.

In this paper, we construct free λ -differential associative algebras with multiple operators and give the Composition-Diamond lemma for such algebras. As applications, we obtain Gröbner-Shirshov bases of free λ -differential Rota-Baxter algebras. Then, linear bases of free λ -differential Rota-Baxter algebras are obtained, which is the same that obtained by Bokut, Chen and Qiu [5], and similar to those obtained by using other methods in the paper of Guo and Keigher [15].

2 Free λ -differential associative algebras with multiple operators

Let K be a field and $\lambda \in K$. A λ -differential associative algebra with multiple operators is a λ -differential algebra R with a set Ω of multi-linear operators. In order to construct the free λ -differential associative algebras with multiple operators, we recall the construction

of free λ -differential algebras firstly. For the detail of the free λ -differential algebras, see [15, 5, 16].

Let X be a set, D a symbol and

$$D^\omega(X) = \{D^i(x) | i \geq 0, x \in X\}$$

where $D^0(x) = x$.

Let $S(D^\omega(X))$ be the free semigroup generated by $D^\omega(X)$. Denote $KS(D^\omega(X))$ the semigroup algebra over the semigroup $S(D^\omega(X))$. Extend linearly $D : KS(D^\omega(X)) \rightarrow KS(D^\omega(X))$ in the following way: for any $u = u_1 u_2 \cdots u_t \in S(D^\omega(X))$, where $u_i \in D^\omega(X)$, define $D(u)$ by induction on t :

If $t = 1$, i.e., $u = D^i(x)$ for some $i \geq 0, x \in X$, then

$$D(u) = D^{i+1}(x).$$

If $t > 1$, then

$$D(u) = \lambda D(u_1) D(u_2 \cdots u_t) + D(u_1)(u_2 \cdots u_t) + u_1 D(u_2 \cdots u_t).$$

Theorem 2.1 ([5, 15, 16]) *$(KS(D^\omega(X)), D)$ is a free λ -differential algebra on X .* ■

Secondly, we construct free λ -differential associative algebras with multiple operators.

Let

$$\Omega = \bigcup_{n=1}^{\infty} \Omega_n$$

where Ω_n is the set of n -ary operators, $\text{ary}(\delta) = n$ if $\delta \in \Omega_n$.

Define

$$\begin{aligned} \mathfrak{L}_0 &= S(D^\omega(X_0)), \quad X_0 = X \\ \mathfrak{L}_1 &= S(D^\omega(X_1)), \quad X_1 = X \cup \Omega(\mathfrak{L}_0) \end{aligned}$$

where

$$\Omega(\mathfrak{L}_0) = \bigcup_{t=1}^{\infty} \{\delta(y_1, y_2, \dots, y_t) | \delta \in \Omega_t, y_i \in \mathfrak{L}_0, i = 1, 2, \dots, t\}.$$

For $n > 1$, define

$$\mathfrak{L}_n = S(D^\omega(X_n)), \quad X_n = X \cup \Omega(\mathfrak{L}_{n-1})$$

where

$$\Omega(\mathfrak{L}_{n-1}) = \bigcup_{t=1}^{\infty} \{\delta(y_1, y_2, \dots, y_t) | \delta \in \Omega_t, y_i \in \mathfrak{L}_{n-1}, i = 1, 2, \dots, t\}.$$

Then we have

$$\mathfrak{L}_0 \subset \mathfrak{L}_1 \subset \cdots \subset \mathfrak{L}_n \subset \cdots.$$

Let

$$\mathfrak{L}(X) = \bigcup_{n \geq 0} \mathfrak{L}_n.$$

Then, it is easy to see that $\mathfrak{L}(X)$ is a semigroup such that $\Omega(\mathfrak{L}(X)) \subseteq \mathfrak{L}(X)$.

For any $v \in \mathfrak{L}(X)$, v has a unique expression

$$v = v_1 v_2 \cdots v_n$$

where each $v_i \in D^\omega(X \cup \Omega(\mathfrak{L}(X)))$. If this is the case, then we define $\text{bre}(v) = n$.

Let $D\langle X; \Omega \rangle = K\mathfrak{L}(X)$ be the semigroup algebra over $\mathfrak{L}(X)$. Then, similar to Theorem 2.1, $D\langle X; \Omega \rangle$ is a λ -differential algebra.


For any $\theta \in \Omega_n$, define

$$\theta : \mathfrak{L}(X)^n \rightarrow \mathfrak{L}(X), (v_1, v_2, \dots, v_n) \mapsto \theta(v_1, v_2, \dots, v_n)$$

and extend linearly $\theta : D\langle X; \Omega \rangle^n \rightarrow D\langle X; \Omega \rangle$. Then it is easy to see that $D\langle X; \Omega \rangle$ is a free λ -differential associative algebra on X with multiple operators Ω .

Now, we describe the elements in $\mathfrak{L}(X)$ by labeled reduced planar rooted forests, see [9, 12].

For any $x \in X_0$, x can be described by a labeled reduced planar rooted tree $\bullet_x$. Then for any element $v = D^i(x)$, $i \geq 1$ can be described by




Then for $u = u_1 u_2 \dots u_n \in S(D^\omega(X))$, we describe u by

$$\bullet_{u_1} \sqcup \bullet_{u_2} \sqcup \cdots \sqcup \bullet_{u_n}$$


which is called a labeled reduced planar rooted forest. For example, if $u = D^3(x_1)D^2(x_2)$, then u can be described by



For any $u \in \Omega(\mathfrak{L}_0)$, say $u = \theta_n(u_1, \dots, u_n)$ where each $u_i \in \mathfrak{L}_0$, u can be described by



For example, if $u = \theta_3(u_1, u_2, u_3) \in \Omega(\mathfrak{L}_0)$ where $u_1 = D(x_1)D^2(x_2)$, $u_2 = D^2(x_3)$, $u_3 = D^3(x_4)D^3(x_5)$, then



Now, for any $v = v_1 v_2 \cdots v_n \in \mathfrak{L}_1 = S(D^\omega(X \cup \Omega(\mathfrak{L}_0)))$ ($v_i \in D^\omega(X \cup \Omega(\mathfrak{L}_0))$, $i = 1, \dots, n$), v can be described by labeled reduced planar rooted forest:

$$\bullet_{v_1} \sqcup \bullet_{v_2} \sqcup \cdots \sqcup \bullet_{v_n}.$$

With the above way, we can describe all the elements in $\mathfrak{S}(X)$ by labeled reduced planar rooted forests.

Therefore, each element in $\mathfrak{S}(X)$ corresponds uniquely to a labeled reduced planar rooted forest.

For $1 \leq t \leq n$, denote by

$$\Theta_n^t = \{(i_1, \dots, i_n) \in \{0, 1\}^n \mid i_1 + \cdots + i_n = t\}.$$

The following proposition is useful in the next section.

Proposition 2.2 *Let $v_t \in D^\omega(X \cup \Omega(\mathfrak{L}(X)))$, $t = 1, \dots, n$. Then*

$$\begin{aligned} D(v_1 v_2 \cdots v_n) &= \sum_{(i_1, \dots, i_n) \in \Theta_n^1} D^{i_1}(v_1) \cdots D^{i_n}(v_n) \\ &\quad + \sum_{t=2}^n \sum_{(i_1, \dots, i_n) \in \Theta_n^t} \lambda^{t-1} D^{i_1}(v_1) \cdots D^{i_n}(v_n). \end{aligned}$$

Proof: If $\lambda = 0$, the result is clear. Assume that $\lambda \neq 0$. Induction on n . For $n = 1, 2$, the result is clear. Now we assume that for $n - 1$ the result is true. Since

$$D(v_1 v_2 \cdots v_n) = \lambda D(v_1) D(v_2 \cdots v_n) + D(v_1) v_2 \cdots v_n + v_1 D(v_2 \cdots v_n)$$

and by induction, we have

$$\begin{aligned} D(v_1 v_2 \cdots v_n) &= \sum_{t=2}^n \sum_{(1, i_2, \dots, i_n) \in \Theta_n^t} \lambda^{t-1} D(v_1) D^{i_2}(v_2) \cdots D^{i_n}(v_n) \\ &\quad + D(v_1) D^0(v_2) \cdots D^0(v_n) \\ &\quad + \sum_{t=1}^{n-1} \sum_{(0, i_2, \dots, i_n) \in \Theta_n^t} \lambda^{t-1} D^0(v_1) D^{i_2}(v_2) \cdots D^{i_n}(v_n) \\ &= \lambda^{n-1} D(v_1) \cdots D(v_n) \\ &\quad + \sum_{t=1}^{n-1} \left(\sum_{(1, i_2, \dots, i_n) \in \Theta_n^t} \lambda^{t-1} D(v_1) D^{i_2}(v_2) \cdots D^{i_n}(v_n) \right. \\ &\quad \left. + \sum_{(0, i_2, \dots, i_n) \in \Theta_n^t} \lambda^{t-1} D^0(v_1) D^{i_2}(v_2) \cdots D^{i_n}(v_n) \right) \\ &= \sum_{t=1}^n \sum_{(i_1, \dots, i_n) \in \Theta_n^t} \lambda^{t-1} D^{i_1}(v_1) \cdots D^{i_n}(v_n). \end{aligned} \quad \blacksquare$$

3 Composition-Diamond lemma for λ -differential associative algebras with multiple operators

In this section, we introduce the notions of Gröbner-Shirshov bases for λ -differential associative algebras with multiple operators and establish the Composition-Diamond lemma for such algebras.

Let X be a set and $D\langle X; \Omega \rangle$ the free λ -differential associative algebra with multiple operators Ω . For any $u \in \mathfrak{L}(X)$,

$$\text{dep}(u) = \min\{n \mid u \in \mathfrak{L}_n\}$$

is called the depth of u . Let $\deg_\Omega(u)$ be the number of the occurrences of $\theta \in \Omega$ in u , for example, if $u = \theta_t(u_1, \dots, u_t) \in \Omega(\mathfrak{L}(X))$, then $\deg_\Omega(u) = 1 + \sum_{i=1}^t \deg_\Omega(u_i)$.

Let X and Ω be well ordered. We define an ordering $>$ on $\mathfrak{L}(X) = \bigcup_{n \geq 0} \mathfrak{L}_n$ by induction on n .

Suppose that $n = 0$, $\mathfrak{L}_0 = S(D^\omega(X_0))$.

Order $D^\omega(X_0)$ by

$$D^i(u) > D^j(v) \text{ if } (i, u) > (j, v) \text{ lexicographically.}$$

Order $\mathfrak{L}_0 = S(D^\omega(X_0))$. For any $u = u_1 \cdots u_t, v = v_1 \cdots v_s \in \mathfrak{L}_0$, (i.e., $u_i, v_j \in D^\omega(X_0)$), define $u > v$ if

$$(\text{bre}(u), u_1, \dots, u_t) > (\text{bre}(v), v_1, \dots, v_s) \text{ lexicographically.}$$

Suppose that $n > 0$. We order $\mathfrak{L}_n$ in three steps.

Firstly, order $X_n = X \cup \Omega(\mathfrak{L}_{n-1})$ by

- (i) $x < u$ if $u \in \Omega(\mathfrak{L}_{n-1})$ and $x \in X$;
- (ii) for $u = \theta_t(u_1, \dots, u_t), v = \delta_s(v_1, \dots, v_s) \in \Omega(\mathfrak{L}_{n-1})$, $u > v$ if

$$(\deg_\Omega(u), \theta_t, u_1, \dots, u_t) > (\deg_\Omega(v), \delta_s, v_1, \dots, v_s) \text{ lexicographically.}$$

Secondly, order $D^\omega(X_n)$. For any $u, v \in X_n$, define

$$D^i(u) > D^j(v) \text{ if } (\deg_\Omega(u), i, u) > (\deg_\Omega(v), j, v) \text{ lexicographically.}$$

Thirdly, order $S(D^\omega(X_n))$. For any $u = u_1 \cdots u_t, v = v_1 \cdots v_s \in S(D^\omega(X_n))$, define $u > v$ if

$$(\deg_\Omega(u), \text{bre}(u), u_1, \dots, u_t) > (\deg_\Omega(v), \text{bre}(v), v_1, \dots, v_s) \text{ lexicographically.}$$

It is easy to check that $>$ is a well ordering on $\mathfrak{L}(X)$. Throughout this paper, this ordering will be used.

Then for any $0 \neq f \in D\langle X; \Omega \rangle$, f has a leading term $\bar{f}$ and

$$f = \alpha_1 \bar{f} + \sum_{i=2}^m \alpha_i u_i$$

where $\bar{f} > u_i, i = 2, \dots, m$ and $0 \neq \alpha_i \in K$. Denote by $lc(f)$ the coefficient of the leading term $\bar{f}$. If $lc(f)$ equals 1, we call f monic.

The proof of the following lemma is straightforward.

Lemma 3.1 Let $u, v \in \mathfrak{L}(X)$, $u > v$, $\theta_n \in \Omega_n$ and $w_j \in \mathfrak{L}(X)$, $j = 1, \dots, n$. Then,

(i) for any i , $1 \leq i \leq n$,

$$\theta_n(w_1, \dots, w_{i-1}, u, w_{i+1}, \dots, w_n) > \theta_n(w_1, \dots, w_{i-1}, v, w_{i+1}, \dots, w_n);$$

(ii) for any $a, b \in \mathfrak{L}(X)$,

$$au > av, \quad ub > vb.$$

■

Lemma 3.2 Let $v_t \in D^\omega(X \cup \Omega(\mathfrak{L}(X)))$, $t = 1, \dots, n$.

(i) If $\lambda \neq 0$, then

$$\overline{D^i(v_1 v_2 \cdots v_n)} = D^i(v_1) D^i(v_2) \cdots D^i(v_n)$$

and the coefficient of $\overline{D^i(v_1 v_2 \cdots v_n)}$ is $\lambda^{(n-1)i}$.

(ii) If $\lambda = 0$, then

$$\overline{D^i(v_1 v_2 \cdots v_n)} = D^i(v_1) v_2 \cdots v_n$$

and the coefficient of $\overline{D^i(v_1 v_2 \cdots v_n)}$ is 1.

It follows that if $u, v \in \mathfrak{L}(X)$ and $u > v$, then $\overline{D(u)} > \overline{D(v)}$.

Proof: Clearly, by using induction on i , (ii) follows from Proposition 2.2. Now, we prove (i).

In Proposition 2.2, we have

$$D(v_1 v_2 \cdots v_n) = \lambda^{n-1} D(v_1) \cdots D(v_n) + \sum_{t=1}^{n-1} \sum_{(i_1, \dots, i_n) \in \Theta_n^t} \lambda^{t-1} D^{i_1}(v_1) \cdots D^{i_n}(v_n) \quad (1)$$

Note that in (1), $D(v_1) \cdots D(v_n) > D^{i_1}(v_1) \cdots D^{i_n}(v_n)$.

We prove the result by induction on i . For $i = 0$ or 1, the result is clear.

Now we assume that the result is true for $i-1$, $i \geq 2$. Since

$$D^i(v_1 v_2 \cdots v_n) = D(D^{i-1}(v_1 v_2 \cdots v_n)),$$

by (1) and by induction, we have

$$\overline{D^i(v_1 v_2 \cdots v_n)} = D^i(v_1) D^i(v_2) \cdots D^i(v_n)$$

and the coefficient of $\overline{D^i(v_1 v_2 \cdots v_n)}$ is $\lambda^{(n-1)i}$. ■

Let $D\langle X; \Omega \rangle$ be a free λ -differential associative algebra with multiple operators Ω on X and $\star \notin X$. By a $\star$ - Ω -word we mean any expression in $\mathfrak{L}(X \cup \{\star\})$ with only one occurrence of $\star$. The set of all $\star$ - Ω -words on X is denoted by $\mathfrak{L}^\star(X)$.


Let u be a $\star$ - Ω -word and $s \in D\langle X; \Omega \rangle$. Then we call

$$u|_s = u|_{\star \mapsto s}$$

an s - Ω -word.

In other words, an s - Ω -word $u|_{\star \mapsto s}$ means that we have replaced the $\star$ of u by s .

For example, if $u = \theta_3(D(x_1)D^2(x_2), D^2(\star), D^3(x_4)D^3(x_5))$, then



Similarly, we can define $(\star_1, \star_2)$ - Ω -words as expressions in $\mathfrak{L}(X \cup \{\star_1, \star_2\})$ with only one occurrence of $\star_1$ and only one occurrence of $\star_2$. Let us denote by $\mathfrak{L}^{\star_1, \star_2}(X)$ the set of all $(\star_1, \star_2)$ - Ω -words. Let $u \in \mathfrak{L}^{\star_1, \star_2}(X)$. Then we call

$$u|_{s_1, s_2} = u|_{\star_1 \mapsto s_1, \star_2 \mapsto s_2}$$

an s_1 - s_2 - Ω -word.

If $\overline{u|_s} = u|_{\bar{s}}$, then we call the s - Ω -word $u|_s$ a normal s - Ω -word.

Note that not each s - Ω -word is a normal s - Ω -word, for example, if

$$u = \theta_3(D(x_1)D^2(\star), D^2(x_3), D^3(x_4)D^3(x_5)) \in \mathfrak{L}^*(X)$$

and $s = xy + 1$, then $u|_s$ is not a normal s - Ω -word. However, if we take

$$u' = \theta_3(D(x_1)\star, D^2(x_3), D^3(x_4)D^3(x_5)) \in \mathfrak{L}^*(X),$$

then $u|_s = u'|_{\star \mapsto D^2(s)}$ and $u'|_{\star \mapsto D^2(s)}$ is a normal $D^2(s)$ - Ω -word. By this way, we can easily prove the following lemma.

Lemma 3.3 *For any s -word $u|_s$, there exist $i \geq 0$ and $u' \in \mathfrak{L}^*(X)$ such that $u|_s = u'|_{D^i(s)}$ and $u'|_{D^i(s)}$ is a normal $D^i(s)$ -word. $\blacksquare$*

By Lemmas 3.1, 3.2, we have

Lemma 3.4 *For any $u, v \in \mathfrak{L}(X), w \in \mathfrak{L}^*(X)$, $u > v \Rightarrow \overline{w|_u} > \overline{w|_v}$. $\blacksquare$*

Let $f, g \in D\langle X; \Omega \rangle$ be monic. Then, there are two kinds of compositions.

- (i) If there exists a $w = \overline{D^i(f)a} = \overline{bD^j(g)}$ for some $a, b \in \mathfrak{L}(X)$ such that $bre(w) < bre(\bar{f}) + bre(\bar{g})$, then we call $(f, g)_w = lc(D^i(f))^{-1}D^i(f)a - lc(D^j(g))^{-1}bD^j(g)$ the intersection composition of f and g with respect to w .
- (ii) If there exists a $u \in \mathfrak{L}^*(X)$ such that $w = \overline{D^i(f)} = u|_{\overline{D^j(g)}}$, then we call $(f, g)_w = lc(D^i(f))^{-1}D^i(f) - lc(D^j(g))^{-1}u|_{D^j(g)}$ the including composition of f and g with respect to w .

Remark: In (i) and (ii), $lc(D^i(f)) = lc(D^j(g)) = 1$ if $\lambda = 0$; $lc(D^i(f)) = \lambda^{(bre(\bar{f})-1)i}$ and $lc(D^j(g)) = \lambda^{(bre(\bar{g})-1)j}$ if $\lambda \neq 0$.

In the above definition, w is called the ambiguity of the composition. Clearly,

$$(f, g)_w \in Id(f, g)$$

where $Id(f, g)$ is the ideal of $D\langle X; \Omega \rangle$ generated by f, g .

It is noted that by an ideal I of $D\langle X; \Omega \rangle$ we mean I is an ideal of $D\langle X; \Omega \rangle$ as associative algebra and closed under D and Ω .

By Lemma 3.4, we have

$$\overline{(f, g)_w} < w.$$

Let $f, g \in D\langle X; \Omega \rangle$ with $\bar{f} = u|_{\overline{D^i(g)}}$ for some $u \in \mathfrak{L}^*(X)$. Then the transformation

$$f \rightarrow f - lc(f)u|_{lc(D^i(g))^{-1}D^i(g)}$$

is called the elimination of the leading word (ELW) of f by g .

Let S be a monic subset of $D\langle X; \Omega \rangle$. Then the composition $(f, g)_w$ is called trivial modulo (S, w) if

$$(f, g)_w = \sum \alpha_i u_i|_{D^{l_i}(s_i)}$$

where each $\alpha_i \in K$, $u_i \in \mathfrak{L}^*(X)$, $s_i \in S$, $u_i|_{D^{l_i}(s_i)}$ is a normal $D^{l_i}(s_i)$ -word and $u_i|_{\overline{D^{l_i}(s_i)}} < w$. If this is the case, we write

$$(f, g)_w \equiv 0 \mod(S, w).$$

In general, for any two polynomials p and q , $p \equiv q \mod(S, w)$ means that $p - q = \sum \alpha_i u_i|_{D^{l_i}(s_i)}$, where each $\alpha_i \in K$, $u_i \in \mathfrak{L}^*(X)$, $s_i \in S$, $u_i|_{D^{l_i}(s_i)}$ is a normal $D^{l_i}(s_i)$ -word and $u_i|_{\overline{D^{l_i}(s_i)}} < w$.

S is called a Gröbner-Shirshov basis in $D\langle X; \Omega \rangle$ if any composition $(f, g)_w$ of $f, g \in S$ is trivial modulo (S, w) .

Lemma 3.5 *Let S be a Gröbner-Shirshov basis in $D\langle X; \Omega \rangle$, $u_1, u_2 \in \mathfrak{L}^*(X)$ and $s_1, s_2 \in S$. If $w = u_1|_{\overline{D^{l_1}(s_1)}} = u_2|_{\overline{D^{l_2}(s_2)}}$, where each $u_i|_{D^{l_i}(s_i)}$ is a normal $D^{l_i}(s_i)$ -word, $i = 1, 2$, then*

$$u_1|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)} \equiv u_2|_{lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} \mod(S, w).$$

Proof: There are three cases to consider.

(i) $\overline{D^{l_1}(s_1)}$ and $\overline{D^{l_2}(s_2)}$ are disjoint. Then there exists a $(\star_1, \star_2)$ - Ω -word Π such that

$$\Pi|_{\overline{D^{l_1}(s_1)}, \overline{D^{l_2}(s_2)}} = u_1|_{\overline{D^{l_1}(s_1)}} = u_2|_{\overline{D^{l_2}(s_2)}}.$$

Then

$$\begin{aligned} & u_2|_{lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} - u_1|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)} \\ &= \Pi|_{\overline{D^{l_1}(s_1)}, lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} - \Pi|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1), \overline{D^{l_2}(s_2)}} \\ &= -\Pi|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1) - \overline{D^{l_1}(s_1)}, lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} \\ & \quad + \Pi|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1), lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2) - \overline{D^{l_2}(s_2)}}. \end{aligned}$$

Let

$$\begin{aligned} -\Pi|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)-\overline{D^{l_1}(s_1)}}, lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} &= \sum \alpha_{2t} u_{2t}|_{D^{l_2}(s_2)}, \\ \Pi|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1), lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)-\overline{D^{l_2}(s_2)}} &= \sum \alpha_{1l} u_{1l}|_{D^{l_1}(s_1)} \end{aligned}$$

where all $u_{2t}|_{D^{l_2}(s_2)}$ and $u_{1l}|_{D^{l_1}(s_1)}$ are normal $D^{l_2}(s_2)$ - and $D^{l_1}(s_1)$ -words, respectively.

By Lemma 3.4, we have $lc(D^{l_i}(s_i)^{-1})D^{l_i}(s_i) - \overline{D^{l_i}(s_i)} < \overline{D^{l_i}(s_i)}$, $i = 1, 2$. It follows that

$$u_1|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)} \equiv u_2|_{lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} \pmod{(S, w)}.$$

(ii) $\overline{D^{l_1}(s_1)}$ and $\overline{D^{l_2}(s_2)}$ have nonempty intersection but do not include each other. Without loss of generality we can assume that $\overline{D^{l_1}(s_1)}a = b\overline{D^{l_2}(s_2)}$ for some $a, b \in \mathfrak{L}(X)$. Then there exists a $\Pi \in \mathfrak{L}^*(X)$ such that

$$\Pi|_{\overline{D^{l_1}(s_1)}a} = u_1|_{\overline{D^{l_1}(s_1)}} = u_2|_{\overline{D^{l_2}(s_2)}} = \Pi|_{b\overline{D^{l_2}(s_2)}}$$

where $\Pi|_{\overline{D^{l_1}(s_1)}a}$ is a normal $D^{l_1}(s_1)a$ -word and $\Pi|_{b\overline{D^{l_2}(s_2)}}$ is a normal $bD^{l_2}(s_2)$ -word. Thus, we have

$$\begin{aligned} &u_2|_{lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} - u_1|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)} \\ &= \Pi|_{lc(D^{l_2}(s_2))^{-1}bD^{l_2}(s_2)} - \Pi|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)a} \\ &= -\Pi|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)a-lc(D^{l_2}(s_2))^{-1}bD^{l_2}(s_2)}. \end{aligned}$$

Since S is a Gröbner-Shirshov basis in $D\langle X; \Omega \rangle$, we have

$$lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)a - lc(D^{l_2}(s_2))^{-1}bD^{l_2}(s_2) = \sum \alpha_j v_j|_{D^{t_j}(s_j)}$$

where each $\alpha_j \in K$, $v_j \in \mathfrak{L}^*(X)$, $s_j \in S$, $v_j|_{\overline{D^{l_1}(s_1)}a} < \overline{D^{l_1}(s_1)}a$ and $v_j|_{D^{t_j}(s_j)}$ is a normal $D^{t_j}(s_j)$ -word. Let $\Pi|_{v_j|_{D^{t_j}(s_j)}} = \Pi_j|_{D^{t_j}(s_j)}$. Then $\Pi_j|_{D^{t_j}(s_j)}$ is also a normal $D^{t_j}(s_j)$ -word. Therefore

$$u_2|_{lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} - u_1|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)} = \sum \alpha_j \Pi_j|_{D^{t_j}(s_j)}$$

with $\Pi_j|_{\overline{D^{t_j}(s_j)}} < w$. It follows that

$$u_1|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)} \equiv u_2|_{lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} \pmod{(S, w)}.$$

(iii) One of $\overline{D^{l_1}(s_1)}$, $\overline{D^{l_2}(s_2)}$ is contained in the other. For example, let $\overline{D^{l_1}(s_1)} = u|_{\overline{D^{l_2}(s_2)}}$ for some $\star$ -word u . Then

$$w = u_2|_{\overline{D^{l_2}(s_2)}} = u_1|_{u|_{\overline{D^{l_2}(s_2)}}} \quad \text{and}$$

$$\begin{aligned} &u_2|_{lc(D^{l_2}(s_2))^{-1}D^{l_2}(s_2)} - u_1|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)} \\ &= u_1|_{lc(D^{l_2}(s_2))^{-1}u|_{\overline{D^{l_2}(s_2)}}} - u_1|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)} \\ &= -u_1|_{lc(D^{l_1}(s_1))^{-1}D^{l_1}(s_1)-lc(D^{l_2}(s_2))^{-1}u|_{\overline{D^{l_2}(s_2)}}}. \end{aligned}$$

Similar to (ii), we can obtain the result. ■

The following theorem is an analogy of Shirshov's Composition lemma for Lie algebras [20], which was specialized to associative algebras by Bokut [4], see also Bergman [3].

Theorem 3.6 (Composition-Diamond lemma) *Let S be a monic subset of $D\langle X; \Omega \rangle$, $Id(S)$ the ideal of $D\langle X; \Omega \rangle$ generated by S and $>$ the order on $\mathfrak{L}(X)$ defined as before. Then the following statements are equivalent:*

- (I) S is a Gröbner-Shirshov basis in $D\langle X; \Omega \rangle$.
- (II) $f \in Id(S) \Rightarrow \bar{f} = u|_{\overline{D^i(s)}}$ for some $u \in \mathfrak{L}^*(X)$, $s \in S$ and $i \geq 0$.
- (III) $Irr(S) = \{w \in \mathfrak{L}(X) | w \neq u|_{\overline{D^i(s)}}, s \in S, u \in \mathfrak{L}^*(X), u|_{D^i(s)} \text{ is a normal } D^i(s)\text{-word}\}$ is a K -basis of $D\langle X; \Omega | S \rangle = D\langle X; \Omega \rangle / Id(S)$.

Proof: (I) $\implies$ (II) Let $0 \neq f \in Id(S)$. Then by Lemma 3.3, we can assume that

$$f = \sum_{i=1}^n \alpha_i u_i |_{lc(D^{l_i}(s_i))^{-1} D^{l_i}(s_i)}$$

where each $\alpha_i \in K$, $u_i \in \mathfrak{L}^*(X)$, $s_i \in S$ and $u_i|_{D^{l_i}(s_i)}$ is a normal $D^{l_i}(s_i)$ -word. Let $w_i = u_i|_{\overline{D^{l_i}(s_i)}}$. We arrange these leading terms in non-increasing order by

$$w_1 = w_2 = \cdots = w_m > w_{m+1} \geq \cdots \geq w_n.$$

Now we prove the result by induction on m .

If $m = 1$, then $\bar{f} = u_1|_{\overline{D^{l_1}(s_1)}}$.

Now we assume that $m \geq 2$. Then $u_1|_{\overline{D^{l_1}(s_1)}} = w_1 = u_2|_{\overline{D^{l_2}(s_2)}}$.

We prove the result by induction on w_1 . Since S is a Gröbner-Shirshov basis in $D\langle X; \Omega \rangle$, by Lemma 3.5, we have

$$u_2|_{lc(D^{l_2}(s_2))^{-1} D^{l_2}(s_2)} - u_1|_{lc(D^{l_1}(s_1))^{-1} D^{l_1}(s_1)} = \sum \beta_j v_j |_{D^{t_j}(s_j)}$$

where $\beta_j \in K$, $s_j \in S$, $v_j \in \mathfrak{L}^*(X)$, $v_j|_{\overline{D^{t_j}(s_j)}} < w_1$ and $v_j|_{D^{t_j}(s_j)}$ is a normal $D^{t_j}(s_j)$ -word. Therefore, since

$$\begin{aligned} & \alpha_1 u_1 |_{lc(D^{l_1}(s_1))^{-1} D^{l_1}(s_1)} + \alpha_2 u_2 |_{lc(D^{l_2}(s_2))^{-1} D^{l_2}(s_2)} \\ &= (\alpha_1 + \alpha_2) u_1 |_{lc(D^{l_1}(s_1))^{-1} D^{l_1}(s_1)} + \alpha_2 (u_2 |_{lc(D^{l_2}(s_2))^{-1} D^{l_2}(s_2)} - u_1 |_{lc(D^{l_1}(s_1))^{-1} D^{l_1}(s_1)}), \end{aligned}$$

we have

$$f = (\alpha_1 + \alpha_2) u_1 |_{lc(D^{l_1}(s_1))^{-1} D^{l_1}(s_1)} + \sum \alpha_2 \beta_j v_j |_{D^{t_j}(s_j)} + \sum_{i \geq 3} \alpha_i u_i |_{lc(D^{l_i}(s_i))^{-1} D^{l_i}(s_i)}.$$

If either $m > 2$ or $\alpha_1 + \alpha_2 \neq 0$, then the result follows from induction on m . If $m = 2$ and $\alpha_1 + \alpha_2 = 0$, then the result follows from induction on w_1 .

(II) $\implies$ (III) For any $f \in D\langle X; \Omega \rangle$, by induction on $\bar{f}$, we have

$$f = \sum_{u_i \in Irr(S), u_i \leq \bar{f}} \alpha_i u_i + \sum_{s_j \in S, v_j |_{\overline{D^{t_j}(s_j)}} \leq \bar{f}} \beta_j v_j |_{D^{t_j}(s_j)}$$

where $\alpha_i, \beta_j \in K$, $v_j|_{D^{l_j}(s_j)}$ is a normal $D^{l_j}(s_j)$ -word. Then $f + Id(S)$ can be expressed by the elements of $Irr(S)$. Now suppose that $\alpha_1 u_1 + \alpha_2 u_2 + \cdots + \alpha_n u_n = 0$ in $D\langle X; \Omega | S \rangle$ with each $0 \neq \alpha_i \in K$, $u_i \in Irr(S)$, Then, in $D\langle X; \Omega \rangle$,

$$g = \alpha_1 u_1 + \alpha_2 u_2 + \cdots + \alpha_n u_n \in Id(S).$$

By (II), we have $\bar{g} = u_i \notin Irr(S)$ for some $1 \leq i \leq n$, a contradiction. Hence, $Irr(S)$ is K -linearly independent. This shows that $Irr(S)$ is a K -basis of $D\langle X; \Omega | S \rangle$.

(III) $\implies$ (I) For any composition $(f, g)_w$ in S , since $(f, g)_w \in Id(S)$ and (III), we have

$$(f, g)_w = \sum \beta_j v_j|_{D^{l_j}(s_j)}$$

where each $\beta_j \in K$, $v_j \in \mathfrak{L}^*(X)$, $s_j \in S$ and $v_j|_{D^{l_j}(s_j)} \leq \overline{(f, g)_w} < w$. This shows (I). $\blacksquare$

4 Gröbner-Shirshov bases for free λ -differential Rota-Baxter algebras

In this section, we obtain a Gröbner-Shirshov basis and a linear basis for a free λ -differential Rota-Baxter algebra. Consequently, we construct the free λ -differential Rota-Baxter algebra on set X .

Let K be a field and $\lambda \in K$. A differential Rota-Baxter algebra of weight λ ([15]), called also λ -differential Rota-Baxter algebra, is an associative K -algebra R with two K -linear operators $P, D : R \rightarrow R$ such that for any $u, v \in R$,

- (i) (Rota-Baxter relation) $P(u)P(v) = P(uP(v)) + P(P(u)v) + \lambda P(uv)$;
- (ii) (λ -differential relation) $D(uv) = \lambda D(u)D(v) + D(u)v + uD(v)$;
- (iii) $D(P(u)) = u$.

Hence, any λ -differential Rota-Baxter algebra is a λ -differential associative algebra with operator $\Omega = \{P\}$.

Throughout this section, we assume $\Omega = \{P\}$ and $D\langle X; \Omega \rangle$ the free λ -differential associative algebra with operator Ω .

Let S be the subset of $D\langle X; \Omega \rangle$ consisting of the following polynomials:

1. $P(u)P(v) - P(uP(v)) - P(P(u)v) - \lambda P(uv)$, $u, v \in \mathfrak{L}(X)$,
2. $D(P(u)) - u$, $u \in \mathfrak{L}(X)$.

Denoted by

$$f(u, v) = P(u)P(v) - P(uP(v)) - P(P(u)v) - \lambda P(uv), \quad u, v \in \mathfrak{L}(X).$$

Lemma 4.1 *Let $j > 0$.*

(i) *If $\lambda \neq 0$, then $\text{mod}(S, D^j(P(u))D^j(P(v)))$*

$$D^j(f(u, v)) \equiv \lambda^j (D^j(P(u))D^j(P(v)) - D^{j-1}(u)D^{j-1}(v)).$$

(ii) *If $\lambda = 0$, then $\text{mod}(S, D^j(P(u))P(v))$*

$$D^j(f(u, v)) \equiv D^j(P(u))P(v) - D^{j-1}(u)P(v).$$

Proof: We just prove (i). The proof of (ii) is similar to (i).

Induction on j . For $j = 1$, we have

$$\begin{aligned}
& D(f(u, v)) \\
&= D(P(u)P(v)) - D(P(uP(v))) - D(P(P(u)v)) - \lambda D(P(uv)) \\
&= \lambda D(P(u))D(P(v)) + D(P(u))P(v) + P(u)D(P(v)) + \\
&\quad - D(P(uP(v))) - D(P(P(u)v)) - \lambda D(P(uv)) \\
&\equiv \lambda D(P(u))D(P(v)) + uP(v) + P(u)v - uP(v) - P(u)v - \lambda uv \\
&\equiv \lambda(D(P(u))D(P(v)) - uv) \pmod{S, D(P(u))D(P(v))}.
\end{aligned}$$

Now we assume that the result is true for $j - 1, j \geq 1$, i.e.,

$$D^{j-1}(f(u, v)) = \lambda^{j-1}(D^{j-1}(P(u))D^{j-1}(P(v)) - D^{j-2}(u)D^{j-2}(v)) + \sum \alpha_i u_i|_{D^{t_i}(s_i)}$$

where each $\alpha_i \in K$, $s_i \in S, u_i|_{\overline{D^{t_i}(s_i)}} < D^{j-1}(P(u))D^{j-1}(P(v))$ and $u_i|_{D^{t_i}(s_i)}$ is a normal $D^{t_i}(s_i)$ -word. Therefore

$$D(\sum \alpha_i u_i|_{D^{t_i}(s_i)}) = \sum \beta_l v_l|_{D^{k_l}(s_l)}$$

where each $v_l|_{D^{k_l}(s_l)}$ is normal $D^{k_l}(s_l)$ -word and $v_l|_{\overline{D^{k_l}(s_l)}} < \overline{D(D^{j-1}(P(u))D^{j-1}(P(v)))} = D^j(P(u))D^j(P(v))$ by Lemma 3.2. So, $\pmod{S, D^j(P(u))D^j(P(v))}$, we have

$$\begin{aligned}
& D^j(f(u, v)) = D(D^{j-1}(f(u, v))) \\
&\equiv \lambda^{j-1}D(D^{j-1}(P(u))D^{j-1}(P(v)) - D^{j-2}(u)D^{j-2}(v)) \\
&\equiv \lambda^{j-1}(\lambda D^j(P(u))D^j(P(v)) + D^j(P(u))D^{j-1}(P(v)) + D^{j-1}(P(u))D^j(P(v)) \\
&\quad - \lambda D^{j-1}(u)D^{j-1}(v) + D^{j-1}(u)D^{j-2}(v) + D^{j-2}(u)D^{j-1}(v)) \\
&\equiv \lambda^{j-1}(\lambda D^j(P(u))D^j(P(v)) + D^{j-1}(u)D^{j-2}(v) + D^{j-2}(u)D^{j-1}(v)) \\
&\quad - \lambda^{j-1}(\lambda D^{j-1}(u)D^{j-1}(v) + D^{j-1}(u)D^{j-2}(v) + D^{j-2}(u)D^{j-1}(v)) \\
&\equiv \lambda^j(D^j(P(u))D^j(P(v)) - D^{j-1}(u)D^{j-1}(v)). \quad \blacksquare
\end{aligned}$$

Theorem 4.2 *With the order $>$ on $\mathfrak{L}(X)$ defined as before, S is a Gröbner-Shirshov basis in $D\langle X; \Omega \rangle$.*

Proof. Denote by $k \wedge l$ the composition of the polynomials of type k and type l . There are two cases $\lambda \neq 0$ and $\lambda = 0$ to consider.

(i) For $\lambda \neq 0$, the ambiguities of all possible compositions of the polynomials in S are list as below:

$$\begin{aligned}
2 \wedge 2 & D^i(D(P(u|_{D^j(D(P(v))}))), \\
2 \wedge 1 & D^i(D(P(u|_{D^j(P(v))D^j(P(w))}))), \\
1 \wedge 2 & D^j(P(u|_{D^i(D(P(v))})))D^j(P(w)), D^j(P(v))D^j(P(u|_{D^i(D(P(w))}))), \\
1 \wedge 1 & D^j(P(u))D^j(P(v))D^j(P(w)), D^j(P(u|_{D^i(P(v))D^i(P(w))})))D^j(P(v')), \\
& D^j(P(v))D^j(P(u|_{D^i(P(w))D^i(P(v'))}))
\end{aligned}$$

where $u \in \mathfrak{L}^*(X)$, $v, w, v' \in \mathfrak{L}(X)$ and $i, j \geq 0$. Now we check that all the compositions are trivial.

$$\begin{aligned}
2 \wedge 2 &= D^i(D(P(u|_{D^j(D(P(v))}))) - u|_{D^j(D(P(v))}) - D^i(D(P(u|_{D^j(D(P(v)-v))}))) \\
&= -D^i(u|_{D^j(D(P(v))}) + D^i(D(P(u|_{D^j(v)}))) \\
&\equiv -D^i(u|_{D^j(v)}) + D^i(u|_{D^j(v)}) \\
&\equiv 0.
\end{aligned}$$

Now, assume that $j > 0$.

$$\begin{aligned}
2 \wedge 1 &= D^i(D(P(u|_{D^j(P(v))D^j(P(w))}))) - D^i(u|_{D^j(P(v))D^j(P(w))}) \\
&\quad - \lambda^{-j} D^i(D(P(u|_{D^j(f(v,w))}))) \\
&\equiv D^i(D(P(u|_{D^j(P(v))D^j(P(w))}))) - D^i(u|_{D^j(P(v))D^j(P(w))}) \\
&\quad - \lambda^{-j} D^i(D(P(u|_{\lambda^j(D^j(P(v))D^j(P(w))-D^{j-1}(v)D^{j-1}(w))}))) \quad (\text{by Lemma 4.1}) \\
&\equiv -D^i(u|_{D^j(P(v))D^j(P(w))}) + D^i(D(P(u|_{D^{j-1}(v)D^{j-1}(w)}))) \\
&\equiv -D^i(u|_{D^{j-1}(v)D^{j-1}(w)}) + D^i(u|_{D^{j-1}(v)D^{j-1}(w)}) \\
&\equiv 0.
\end{aligned}$$

$$\begin{aligned}
1 \wedge 2 &= \lambda^{-j} D^j(f(u|_{D^i(D(P(v))}), w)) - D^j(P(u|_{D^i(D(P(v))-v)})D^j(P(w)) \\
&\equiv \lambda^{-j} (\lambda^j (D^j(P(u|_{D^i(D(P(v))})))D^j(P(w)) - D^{j-1}(u|_{D^i(D(P(v))})D^{j-1}(w))) \\
&\quad - D^j(P(u|_{D^i(D(P(v))})D^j(P(w)) + D^j(P(u|_{D^i(v)})D^j(P(w)) \quad (\text{by Lemma 4.1}) \\
&\equiv 0.
\end{aligned}$$

Similarly, we can prove another case of $1 \wedge 2$ to be also trivial.

$$\begin{aligned}
1 \wedge 1 &= \lambda^{-j} D^j(f(u, v))D^j(P(w)) - \lambda^{-j} D^j(P(u))D^j(f(u, w)) \\
&\equiv (D^j(P(u))D^j(P(v)) - D^{j-1}(u)D^{j-1}(v))D^j(P(w)) \\
&\quad - D^j(P(u))(D^j(P(v))D^j(P(w)) - D^{j-1}(v)D^{j-1}(w)) \quad (\text{by Lemma 4.1}) \\
&\equiv -D^{j-1}(u)D^{j-1}(v))D^{j-1}(w) + D^{j-1}(u)D^{j-1}(v))D^{j-1}(w) \\
&\equiv 0.
\end{aligned}$$

Another two cases in $1 \wedge 1$ can be easily checked.

For $j = 0$, the proof is similar. We omit the details.

(ii) For $\lambda = 0$, the ambiguities of all possible compositions of the polynomials in S are list as below:

$$\begin{aligned}
2 \wedge 2 & D^i(D(P(u|_{D^j(D(P(v))}))), \\
2 \wedge 1 & D^i(D(P(u|_{D^j(P(v))P(w)}))), \\
1 \wedge 2 & D^j(P(u|_{D^i(D(P(v))}))P(w), D^j(P(v))P(u|_{D^i(D(P(w))})), \\
1 \wedge 1 & D^j(P(u))P(v)P(w), D^j(P(u|_{D^i(P(v))P(w)}))P(v'), D^j(P(v))P(u|_{D^i(P(w))P(v')})
\end{aligned}$$

where $u \in \mathfrak{L}^*(X)$, $v, w, v' \in \mathfrak{L}(X)$ and $i, j \geq 0$.

Similar to the proof of (i), we can prove that all compositions are trivial modulo S . ■

Let Y and Z be two subsets of $\mathfrak{L}(X)$. Define the alternating product of Y and Z (see also [11]):

$$\Lambda(Y, Z) = (\cup_{r \geq 1} (YP(Z))^r) \cup (\cup_{r \geq 0} (YP(Z))^r Y) \cup (\cup_{r \geq 1} (P(Z)Y)^r) \cup (\cup_{r \geq 0} (P(Z)Y)^r P(Z)).$$

Define

$$\Phi_0 = S(D^\omega(X)),$$

and for $n > 0$,

$$\Phi_n = \Lambda(\Phi_0, \Phi_{n-1}).$$

Let

$$\Phi(D^\omega(X)) = \bigcup_{n \geq 0} \Phi_n.$$

By Theorem 4.2 and Theorem 3.6, we have the following theorems.

Theorem 4.3 ([5]) *$\text{Irr}(S) = \Phi(D^\omega(X))$ is a K -basis of $D\langle X; \Omega|S \rangle$.*

Theorem 4.4 ([5]) *$D\langle X; \Omega|S \rangle$ is a free λ -differential Rota-Baxter algebra on set X with a linear basis $\Phi(D^\omega(X))$.*

We note here that Theorem 4.4 is also obtained by L. Guo and W. Keigher in [15] by using the planar decorated rooted trees.

Acknowledgement: The authors would like to thank Professor L.A. Bokut for his guidance, useful discussions and enthusiastic encouragement in writing up this paper.

References

- [1] M. Aguiar and M. Moreira, Combinatorics of free Baxter algebra, Electron J. Combin, to appear. [arXiv:math.CO/0510169].
- [2] G. Baxter, An analytic problem whose solution follows from a simple algebraic identity, Pacific J. Math., **10**(1960), 731-742.
- [3] G. M. Bergman, The diamond lemma for ring theory, Adv. Math., **29**(1978), 178-218.
- [4] L. A. Bokut, Imbeddings into simple associative algebras, Algebra i Logika, **15**(1976), 117-142.
- [5] L. A. Bokut, Yuqun Chen and Jianjun Qiu, Gröbner-Shirshov bases for associative algebras with multiple operators and free Rota-Baxter algebras, [arXiv.org/abs/0805.0640].

- [6] B. Buchberger, An algorithm for finding a basis for the residue class ring of a zero-dimensional polynomial ideal [in German], Ph.D. thesis, University of Innsbruck, Austria, (1965).
- [7] B. Buchberger, An algorithmical criteria for the solvability of algebraic systems of equations [in German], *Aequationes Math.*, **4**(1970), 374-383.
- [8] P. Cartier, On the structure of free Baxter algebras, *Adv. in Math.*, **9** (1972), 253-265.
- [9] V. Drensky, R. Holtkamp, Planar trees, free nonassociative algebras, invariants, and elliptic integrals, *Algebra and Discrete Mathematics*, **2**(2008), 1-41.
- [10] K. Ebrahimi-Fard, J. M. Gracia-Bondia and F. Patras, Rota–Baxter algebras and new combinatorial identities, *Lett. Math. Phys.* **81**(2007), 61-65.
- [11] K. Ebrahimi-Fard, L. Guo, Rota–Baxter algebras and dendriform dialgebras, *Journal of Pure and Applied Algebra*, **212**(2)(2008), 320-339.
- [12] K. Ebrahimi-Fard, L. Guo, Free Rota–Baxter algebras and rooted trees, *J. Algebra Applications*, **7**(2008), 167-194.
- [13] L. Guo, W. Keigher, Baxter algebras and Shuffle products, *Adv. Math.*, **150** (2000), 117-149.
- [14] L. Guo, W. Keigher, On free Baxter algebras: completions and the internal construction, *Adv. Math.*, **151**(2000), 101-127.
- [15] L. Guo, W. Keigher, On differential Rota–Baxter algebras, *Journal of Pure and Applied Algebra*, **212**(2008), 522-540.
- [16] E. Kolchin, *Differential algebras and algebraic groups*, Academic Press, NewYork, 1973.
- [17] A. G. Kurosh, Free sums of multiple operator algebras, *Siberian. Math. J.*, **1**(1960), 62-70.
- [18] A. G. Kurosh, Multi operators rings and algebras, *Russian Math. Surveys. Uspekhi*, **24**(1)(1969), 3-15.
- [19] G.-C. Rota, Baxter algebras and combinatorial identities I, *Bull. Amer. Math. Soc.*, **5**(1969), 325-329.
- [20] A. I. Shirshov, Some algorithmic problem for Lie algebras, *Sibirsk. Mat. Z.* **3**(1962), 292-296 (in Russian); English translation in *SIGSAM Bull.*, **33**(2)(1999), 3-6.